

Vaccine Name	Contents	What it protects against	When your child needs it
Pentacel	Dtap-IPV, Hib	diphtheria, tetanus, pertussis, polio(myelitis), and invasive disease due to Haemophilus influenzae type b	2 months, 4 months, 6 months
Prevnar 20	Pneumococcal 20-valent Conjugate	-invasive disease caused by 20 Streptococcus pneumoniae strains (1, 3, 4, 5, 6A, 6B, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F, and 33F) in individuals 6 weeks and older. -otitis media (middle ear infection) caused by 7 of the 20 strains in individuals 6 weeks through 5 years.	2 months, 4 months, 6 months
RotaTeq	Rotavirus (oral)	RotaTeq is an oral vaccine used to help prevent rotavirus infection in children. Rotavirus infection can cause fever, vomiting, and diarrhea that can be severe and can lead to loss of body fluids (dehydration), hospitalization, and even death in some children.	2 months, 4 months, 6 months
Hep B	Hepatitis B	Hepatitis B is a liver infection caused by a virus that can spread through bodily fluids	birth, 1 month, 9 months
MMR	MMR	Measles: Causes an uncomfortable rash and high fever Mumps: Causes a fever and swollen glands under the ears or jaw Rubella: Causes a rash and fever, and can be passed on to a pregnant person, who could develop serious birth defects or lose the pregnancy	12 months
Varicella	Varicella	chickenpox	12 months
Hep A	Hepatitis A	hepatitis A, a serious liver disease	12 months
ProQuad	MMRV	Measles, Mumps, Rubella, Varicella	4 years
Quadracel	Dtap-IPV	diphtheria, tetanus, pertussis, and polio(myelitis); similar to Pentacel but without Hib	4 years
Tdap	Tdap	tetanus, diphtheria, and pertussis (whooping cough)	11 years+
Menquadfi	MenACWY	invasive meningococcal disease caused by Neisseria meningitidis serogroups A, C, W, and Y	11 years, 16 years
Gardasil	HPV	human papillomavirus HPV types 6, 11, 16, 18, 31, 33, 45, 52, and 58, including: Cervical, vaginal, and vulvar cancers in females Anal cancer Certain head and neck cancers, such as throat and back of mouth cancers Genital warts in both males and females	11 years, 12-15 years
Bexsero	Men B	meningococcal group B disease (also known as meningitis B) caused by Neisseria meningitidis bacteria	16 years